

2023 CLABSI External Validation Toolkit

The 2023 CLABSI External Validation Toolkit is a HAI-specific supplement to be used in conjunction with the 2023 NHSN Patient Safety External Validation Toolkit (2023 PS EVT). It is intended to help guide the external validation process specifically for CLABSI with step-by-step instructions and screenshots from NHSN.

Table of Contents

Section 1. Facility Selection	2
1.1 Steps Applicable to all Facility Selection Methods	2
Generate Datasets and Modify Report.....	2
1.2 Calculate Ranking and Selection:.....	6
Method 1: Prioritizing Facilities with Highest Likelihood of Event Occurrence	6
Method 2: Cumulative Attributable Difference (CAD) Approach.....	7
Method 3: Stratified Random Sampling	10
Section 2. Download (“freeze”) the facility’s reported data from NHSN	14
Section 3. Notify facilities of the planned validation and request the required laboratory line listings	15
Section 4. Develop the medical record sampling frame for each selected facility	15
Section 5. Medical Record Selection	16
Section 6. Site Visit Activities.....	18
6.1 Structured Medical Records Review.....	18
6.2 Review risk adjustment variables	18
6.3 Review denominator collection methods and documentation.....	18
6.4 (Optional) Template for Central Line-Associated Bloodstream Infection (CLABSI) Validation Discrepancies Discussion with Facilities	20
6.5 Documentation of Electronic CLABSI Denominator Validation.....	21
6.6 Contact Information for Manual CLABSI Denominator Validation	22
6.7 CLABSI Denominator Counting Survey (with Key)	23
6.8 (Optional) 2023 CLABSI Validation Summary	28

Section 1. Facility Selection


1.1 Steps Applicable to all Facility Selection Methods

Generate Datasets and Modify Report


1. Generate new datasets in NHSN to ensure any data updates are included for analysis. On the NHSN Landing Page, navigate to Patient Safety Component → [YOUR State/Jurisdiction Users' Group]. Select the Analysis tab and click Generate Data Sets. For Beginning, enter 01/2023 and for Ending, 12/2023 (or other dates corresponding to the timeframe being validated) for the data set time period. Click the Generate New button. Allow the dataset generation process to complete; you can leave NHSN during the generation process.

The screenshot displays the NHSN - National Healthcare Safety Network interface. On the left is a sidebar with the following links: NHSN Home, Alerts, Dashboard, Reporting Plan, Patient, Event, Procedure, Summary Data, COVID-19, Import/Export, Surveys, Analysis, Users, Facility, Group, and Logout. The main content area is titled 'Generate Data Sets (Patient Safety)'. Below this title is a 'Reporting Data Sets' tab. A gear icon is present next to the text 'Include data for the following time period:'. Below this text are two date pickers: 'Beginning' (01/2017) and 'Ending' (03/2021). A 'Clear Time Period' button is located to the right of the date pickers. At the bottom left of the main content area is a 'Generate Reporting Data Sets' button. At the bottom right is a yellow box containing the text: 'Last Generated: February 24, 2021 3:31 PM to include data beginning 01/2017 and ending 09/2020'.

2. After successful dataset generation, navigate to Analysis → Reports to display the tree view list of all analysis reports available within NHSN's analysis tool.



3. Use the tree view structure to select HAI Risk Adjusted Measure Reports, 2015 Baseline (Baseline Set 2), CLABSI and MBI-LCBI, SIR – ACH CLABSI Data. If you are validating Critical Access Hospitals (CAHs), Inpatient Rehab Facilities (IRFs), or Long-term Acute Care Hospitals (LTACHs), select the SIR report that corresponds with that facility type. Click the Modify Report button to proceed to the modification window.



4. In the modification window, there are two key areas to modify, one that controls the time interval of data that are analyzed and displayed and one that controls the level of aggregation of those data.
 - 4a. Under Title/Format tab, select xls format. Then navigate to the Time Period tab to define the time period of data that is included in the report to be exported. Set Date Variable to "summaryYr," Beginning and Ending to 2023, or to the year of data being validated.

Modify "SIR - ACH CLABSI Data (2015 Baseline)"

☐ Show descriptive variable names ([Print List](#))

Title/Format	Time Period	Filters	Display Options								
Time Period: <table> <tr> <th>Date Variable</th> <th>Beginning</th> <th>Ending</th> <th></th> </tr> <tr> <td>summaryYr ▼</td> <td>2023</td> <td>2023</td> <td> Clear Time Period</td> </tr> </table> <p><input type="checkbox"/> Enter Date variable/Time period at the time you click the Run button</p>				Date Variable	Beginning	Ending		summaryYr ▼	2023	2023	Clear Time Period
Date Variable	Beginning	Ending									
summaryYr ▼	2023	2023	Clear Time Period								

4b. Navigate to the "Filters" tab. In the row of drop-down boxes, select "bsiPlan," "equal," and enter "Y."

Title/Format	Time Period	Filters	Display Options												
Additional Filters: Show Clear															
<table> <tr> <td>AND OR</td> <td colspan="2"></td> <td> Add group</td> </tr> <tr> <td>AND OR</td> <td colspan="2"></td> <td> Add rule</td> </tr> <tr> <td></td> <td>bsiPlan ▼</td> <td>equal ▼</td> <td>Y Delete</td> </tr> </table>				AND OR			Add group	AND OR			Add rule		bsiPlan ▼	equal ▼	Y Delete
AND OR			Add group												
AND OR			Add rule												
	bsiPlan ▼	equal ▼	Y Delete												

4c. Under the Display Options section, use the Group by option to view the data at a particular level of aggregation. Change the Group by option to "summaryYr."

Modify "SIR - Acute Care Hospital Data"					
<input type="checkbox"/> Show descriptive variable names (Print List)					
Title/Format	Time Period	Filters	Display Options		
SIR Options: <table> <tr> <td>Group by:</td> <td>summaryYH ▼</td> </tr> </table> <div> Cumulative summaryYH summaryYM summaryYQ summaryYr </div>				Group by:	summaryYH ▼
Group by:	summaryYH ▼				

- After making the above modifications, scroll to the bottom of the modification window. Click the Export button to export the data selected by your modifications. This will open the “Export Analysis Data Set” window.
- Use the default file format (.csv) and select the bullet “Export Analysis Data Set using Modifications” to export the data. Click the Export button to begin the export process. NHSN will create a .zip file with your SIR data report in it and prompt you to specify a location to save the file on your computer.

Export Analysis Data Set

Analysis Data Set: bs2_CLAB_RatesICU

Export Format: delimited file (comma-separated values) (*.csv) ▼

☐ Export Entire Analysis Data Set
☒ Export Analysis Data Set using Modifications

Export
Cancel

- The exported SIR report will be displayed at several levels of aggregation. Select the orgID level, as illustrated in the screenshot below, to get an unduplicated list of facilities in your jurisdiction.

infCount	numPred	numclday	SIR_pval	SIR	sir95ci	SIR_pctl	locationType	locCDC	orgID	facType	months			
5	2.84691	2423	0.229	1.756	0.643, 3.893	83	SIR for all locations in all facilities in group							
5	0.0422	56					CC	SIR for all location types in all facilities in group						
0	1.55251	905	0.2117	0	1.930	25	CC_N							
0	0.60882	808					CC_ONC							
0	0.00405	6					STEP							
0	0.04232	65					WARD							
0	0.59702	583					WARD_ONC							
5	0.0422	56						IN:ACUTE:CC:C	SIR for each location type in all facilities in group					
0	1.2868	800	0.2762	0	2.328			IN:ACUTE:CC:NURS						
0	0.06326	25						IN:ACUTE:CC:NURS_IV						
0	0.60882	808						IN:ACUTE:CC:ONC_M						
0	0.20245	80						IN:ACUTE:CC_STEP:NURS						
0	0.00405	6						IN:ACUTE:STEP:NURS						
0	0.04232	65						IN:ACUTE:WARD:M						
0	0.59702	583						IN:ACUTE:WARD:ONC_HONC						
5	1.10226	1110	0.0065	4.536	1.662, 10.054	100			10018	HOSP-GEN		*THIS IS THE LEVEL TO EVALUATE*		
0	1.74466	1313	0.1747	0	1.717	5			13860	HOSP-GEN				
5	0.0422	56					CC		10018	HOSP-GEN		Facility-specific SIRs combining all location types		
0	0.40487	175					CC_N		10018	HOSP-GEN				
0	0.60882	808					CC_ONC		10018	HOSP-GEN				
0	0.00405	6					STEP		10018	HOSP-GEN				
0	0.04232	65					WARD		10018	HOSP-GEN				
0	1.14764	730	0.3174	0	2.610	25	CC_N		13860	HOSP-GEN				
0	0.59702	583					WARD_ONC		13860	HOSP-GEN				
5	0.0422	56						IN:ACUTE:CC:C	10018	HOSP-GEN		Facility and location type-specific SIRs		
0	0.13916	70						IN:ACUTE:CC:NURS	10018	HOSP-GEN				
0	0.06326	25						IN:ACUTE:CC:NURS_IV	10018	HOSP-GEN				
0	0.60882	808						IN:ACUTE:CC:ONC_M	10018	HOSP-GEN				
0	0.20245	80						IN:ACUTE:CC_STEP:NURS	10018	HOSP-GEN				
0	0.00405	6						IN:ACUTE:STEP:NURS	10018	HOSP-GEN				
0	0.04232	65						IN:ACUTE:WARD:M	10018	HOSP-GEN				
								IN:ACUTE:CC:NURS	13860	HOSP-GEN				
								IN:ACUTE:WARD:ONC_HONC	13860	HOSP-GEN		Facility and specific location SIRs		
								IN:ACUTE:WARD:M	10018	HOSP-GEN	1			
5	0.0422	56					CC	IN:ACUTE:CC:C	10018	HOSP-GEN	2			
0	0.07589	45					CC_N	IN:ACUTE:CC:NURS	10018	HOSP-GEN	1			

This Excel spreadsheet shows seven different levels of aggregation for the NHSN Facility CLABSI SIR download. Select the tier (shown in black above) that identifies the facility-specific SIR for CLABSI combining all location types.

1.2 Calculate Ranking and Selection:

Method 1: Prioritizing Facilities with Highest Likelihood of Event Occurrence

1. Open the exported SIR report in Excel and select the aggregation level that provides a facility-specific SIR at the orgID level (shown in black in the screenshot seen in Section 1.1 step 7). This will allow you to explore the level of exposure risk for CLABSIs and measured performance for each facility. An easy way to find this level of aggregation is to look at the “loccdc” column and scroll until it is blank. The unique orgIDs will then begin to list in ascending order. Once you see the list go through the highest orgID and start over at the smallest orgID, that is where the unduplicated facility list ends. You will also notice that once the list of orgIDs starts over, the column “locationtype” will begin to have data as well. *Tip: the columns “loccdc” and “locationtype” are **blank** for the rows you want.*
2. Copy this information to a new spreadsheet. Arrange the facilities in descending rank order according to SIR, and create three new columns titled “Delta,” “Stratum,” and “Targeted Selection Number.”
3. Calculate Delta for each facility/row using the formula =ABS[row cell under InfCount]-[row cell under numPred]. Delta will be used only where an SIR is not calculated by NHSN.
4. Select the top tertile (33%) of facilities by predicted number (numPred) of CLABSIs in surveillance locations. This top tertile of facilities where CLABSIs in surveillance locations are most expected and may have the greatest potential for surveillance and prevention impact.
5. Within the top tertile, sort by SIR in descending order, and identify the current median SIR for the top tertile. To sort just the top tertile, highlight the entire row for each facility in the top tertile, and click “Data,” “Sort”; Sort by “Column” (select SIR), “Sort On” (cell values), and “Order” (largest to smallest).
6. Within the top tertile, assign stratum A to facilities with SIR above the current median SIR, stratum B for remaining facilities with SIR less than or equal to the median and above zero, and stratum C for facilities with SIR = zero (but not missing). Note that some facilities will not have a calculated SIR; do not include these in the strata (see step 9 below).
7. Re-sort within each stratum A, B, and C, by numPred from highest to lowest. To sort just one stratum at a time, highlight the entire row for each facility in the first stratum, and click “Data,” “Sort”; Sort by “Column” (select numPred), “Sort On” (cell values), and “Order” (highest to lowest). Repeat this process for the next two strata, one-by-one.
8. Assign sequential Targeted Selection Numbers to facilities by selecting the highest available numPred from each stratum, alternating through strata A, B, and C. For example, the facility with the highest numPred from stratum A would be Targeted Selection Number=1, the facility with the highest numPred from stratum B would be Targeted Selection Number=2, and the facility with the highest numPred from stratum C would be Targeted Selection Number=3. Return to stratum A and assign the facility with the next highest numPred as Targeted Selection Number=4. Continue alternating strata until no facilities remain or the target number of facilities is reached (typically 18 or 21; refer to the 2023 PS EVT for facility sample size recommendations). If additional facilities are needed, repeat steps 4-8 using the second and then third tertile based on risk level.

9. If additional facilities are needed to reach the targeted number after step 8 is complete, sort the remaining facilities without a calculated SIR by Delta in descending order, starting with the highest, and select facilities from the top of the list until targeted number is reached.
10. After the targeted selection is complete, randomly select additional 5% of remaining facilities from ALL tertiles. The targeted facilities along with the 5% randomly selected make up the total sample.

Method 2: Cumulative Attributable Difference (CAD) Approach

1. Open the exported SIR report in Excel and select the aggregation level that provides a facility-specific SIR at the orgID level (shown in black in the screenshot seen in Section 1.1 step 7). This will allow you to explore the level of exposure risk for CLABSIs and measured performance for each facility. An easy way to find this level of aggregation is to look at the “loccdc” column and scroll until it is blank. The unique orgIDs will then begin to list in ascending order. Once you see the list go through the highest orgID and start over at the smallest orgID, that is where the unduplicated facility list ends. You will also notice that once the list of orgIDs starts over, the column “locationtype” will begin to have data as well. *Tip: the columns “loccdc” and “locationtype” are blank for the rows you want.*
2. If there are 30 or fewer facilities in your jurisdiction, stop here and validate them all. If there are more than 30 facilities, proceed through the following steps to create facility sampling frame.
3. Select the rows from the aggregation level being evaluated and copy this information to a new spreadsheet. Insert a row above your data and copy the header row so you can identify the variables on the new spreadsheet. Next, sort the facilities by numPred (number of predicted events) in descending order (high to low).

InfCount	numPred	numclay	SIR_pval	SIR	sir95ci	locationTy	locCDC	orgID	facType
13	18.51959	15267	0.1921	0.702	0.390, 1.170			100008	HOSP-GEN
22	15.32671	9910	0.1034	1.435	0.922, 2.138			100030	HOSP-CHLD
10	9.736101	8387	0.8926	1.027	0.522, 1.831			100001	HOSP-GEN
8	9.542312	7958	0.6509	0.838	0.389, 1.592				
1	9.064373	7682	0.0013	0.11	0.006, 0.544				
2	7.578169	6272	0.0235	0.264	0.044, 0.872				
7	5.689505	4581	0.5585	1.23	0.538, 2.434				
4	5.504663	4879	0.558	0.727	0.251, 1.735				
2	3.159258	2784	0.5651	0.633	0.106, 2.092				
2	2.437844	2304	0.8601	0.82	0.138, 2.710				
0	1.945079	1724	0.143	0	1.540				
3	1.572374	1812	0.2846	1.908	0.485, 5.193				
1	1.527251	1760	0.7659	0.655	0.033, 3.229			100010	HOSP-GEN
2	1.329405	1357	0.5333	1.504	0.252, 4.970			100032	HOSP-GEN
0	1.242188	1101	0.2888	0	2.412			100049	HOSP-GEN
2	1.087298	1253	0.3934	1.839	0.308, 6.077			100005	HOSP-GEN
1	1.052644	933	1	0.95	0.048, 4.685			100007	HOSP-GEN
1	0.915007	934						100040	HOSP-GEN
0	0.745198	989						100026	HOSP-GEN
0	0.719899	823						100004	HOSP-GEN
2	0.669096	888						100023	HOSP-GEN

Sort the facilities in descending order of number of predicted infections (numPred) and compute the 75th percentile value of the variable numPred

4. Identify the 75th percentile of numPred for the validation period (minimum of two quarters of data) using the Percentile.inc function in Excel by clicking on the function button (f_x) to the left of the white text box and selecting “Percentile.inc.” A Function Arguments window will open and require an array and K values. For the “array” argument, select the column of your spreadsheet containing numPred values. For the “K” argument, enter the percentile value to be generated (0.75), making sure to use a decimal. Click OK and the cell where the function was entered will now show the 75th percentile value.

Function Arguments

PERCENTILE.INC

Array: C2:C50

K: 0.75

= PERCENTILE.INC(C2:C50,0.75)

Returns the k-th percentile of values in a range, where k is in the range 0..1, inclusive.

Array is the array or range of data that defines relative standing.

Formula result = PERCENTILE.INC(C2:C50,0.75)

[Help on this function](#)

OK Cancel

- Use the numPred value corresponding to the 75th percentile as the minimum threshold value for selection of facilities eligible for validation. If this value is greater than 1, use the 75th percentile numPred value, otherwise use numPred=1 as the minimum threshold value.

infCount	numPred	numclday	SIR_pval	SIR	sir95ci	locationTy	locCDC	orgID	facType
13	18.51959	15267	0.1921	0.702	0.390, 1.170			100008	HOSP-GE
22	15.32671	9910	0.1034	1.435	0.922, 2.138			100030	HOSP-CH
10	9.736101	8387	0.8926	1.027	0.522, 1.831			100014	HOSP-GE
8	9.542312	7958	0.6509	0.838	0.389, 1.592			100046	HOSP-GE
1	9.064373	7682	0.0013	0.11	0.006, 0.544			100001	HOSP-GE
2	7.578169	6272	0.0235	0.264	0.044, 0.872			100002	HOSP-GE
7	5.689505	4581	0.5585	1.23	0.538, 2.434			100022	HOSP-GE
4	5.504663	4879	0.558	0.727	0.231, 1.753			100027	HOSP-GE
2	3.159258	2784	0.5651	0.633	0.308, 1.000			100005	HOSP-GE
2	2.437844	2304	0.8601	0.82	0.308, 1.000			100005	HOSP-GE
0	1.945079	1724	0.143	0	0.308, 1.000			100005	HOSP-GE
3	1.572374	1812	0.2846	1.908	0.308, 1.000			100005	HOSP-GE
1	1.527251	1760	0.7659	0.655	0.308, 1.000			100005	HOSP-GE
2	1.329405	1357	0.5333	1.504	0.308, 1.000			100005	HOSP-GE
0	1.242188	1101	0.2888	0	0.308, 1.000			100005	HOSP-GE
2	1.087298	1253	0.3934	1.839	0.308, 6.077			100005	HOSP-GE

75th percentile value of numPred = 5.5. Select facilities with numPred > 5.5. Only facilities in the red box (numPred > 5.5) are included in the sampling frame for targeted validation.

- Create a subset of facilities that includes facilities with predicted number of CLABSI events greater than the threshold. In the example above, the 75th percentile value of the numPred variable was 5.5. All facilities with numPred value above 5.5 are selected for inclusion in the validation sampling frame.
- If the sampling frame derived from the 75th percentile of numPred consists of 30 or fewer facilities, select all facilities for validation, plus an additional random sample of 5% of facilities where numPred was less than the 75th percentile value. Refer to Table 1 below for 3 random number generation methods. In the example above, the number of facilities with numPred value > 5.5 is fewer than 30 so all facilities with numPred value > 5.5 are selected for validation and a 5% random sample is selected from the facilities with numPred value ≤ 5.5.
- If sampling frame consists of greater than 30 facilities, select 30 facilities based on the criteria described in section B below.

A. Observed Events

- The Cumulative Attributable Difference (CAD) approach focuses on the difference between the predicted number of CLABSIs and actual observed CLABSIs (reported). The infCount is an aggregated count of observed CLABSIs for individual surveillance locations.
- Create a column titled CAD next to numPred and compute the CAD values for each line by subtracting numPred from infCount (observed – predicted).

B. Facility Selection: use this step if the sampling frame consists of greater than 30 facilities

- 1) Divide the total facilities in the sampling frame into two strata.
 - Create a new column, “stratum,” and assign each facility to either Stratum 1 or Stratum 2:
 - Stratum 1: Includes all facilities in the sampling frame that had zero infCount value, that is, zero reported observed events for the validation time frame.
 - Stratum 2: Includes all facilities in the sampling frame with non-zero infCount value, that is, non-zero reported observed events for the validation time frame.
- 2) Stratum 1 (where facility reported zero events): Filter for Stratum 1 facilities (where infCount=0) and sort by ascending CAD value so stratum 1 facilities with the lowest CAD value are at the top, shown in the green column below. Select the first 15 facilities from Stratum 1.

infCount	numpred	CAD	numclay	SIR_pval	SIR	sir95ci	SIR_pctl	locationTy	locCDC	orgID	facType
0	15.11302	-15.1130	150								HOSP-GEN
0	12.12433	-12.1243	165								HOSP-GEN
0	11.03699	-11.0370	68								HOSP-GEN
0	10.03699	-10.0370	68								HOSP-GEN
0	9.113023	-9.1130	150								HOSP-GEN
0	8.124325	-8.1243	165								HOSP-GEN
0	8.113023	-8.1130	150								HOSP-GEN
0	8.011302	-8.0113	15								HOSP-GEN
0	5.011302	-5.0113	15								HOSP-GEN
0	4.036994	-4.0370	68								HOSP-GEN
0	3.036994	-3.0370	68								HOSP-GEN
0	3.011302	-3.0113	15								HOSP-GEN
0	2.036994	-2.0370	68								HOSP-GEN
0	1.011302	-1.0113	15								HOSP-GEN

- 3) Stratum 2 (facilities with non-zero events): Filter for Stratum 2 facilities (where infCount > 0). Sort the facilities by ascending CAD value so the facilities with the lowest CAD value are at the top, shown in the green column below. Select the first 15 facilities from Stratum 2.

infCount	numPred	CAD	numclday	SIR_pval	SIR	sir95ci	locationTy	locCDC	orgID	facType
1	9.064373	-8.06437	7682	0.0013						HOSP-GEN
2	7.578169	-5.57817	6272	0.0235						HOSP-GEN
13	18.51959	-5.51959	15267	0.1921						HOSP-GEN
8	9.542312	-1.54231	7958	0.6509						HOSP-GEN
4	5.504663	-1.50466	4879	0.558						HOSP-GEN
10	9.736101	0.263899	8387	0.8926						HOSP-GEN
7	5.689505	1.310495	4581	0.5585						HOSP-GEN
22	15.32671	6.673286	9910	0.1034						HOSP-CHLD

Compute the CAD values for all facilities in the sampling frame. Filter for stratum 2 facilities, where infCount > 0. Sort by ascending CAD value (lowest values on the top). If the sampling frame has greater than 15 facilities, select the top 15 facilities.

- 4) If there are insufficient facilities in either of the strata, supplement the sample from other strata to reach the required number of facilities for the validation sample.

Note: Remember to randomly select 5% of remaining facilities with a numPred less than the 75th percentile value.

Method 3: Stratified Random Sampling

1. Open the exported SIR report in Excel and select the aggregation level that provides a facility-specific SIR at the orgID level (shown in black in the screenshot seen in Section 1.1 step 7) so you have an unduplicated list of all facilities reporting data for CLABSI during the validation timeframe. An easy way to find this level of aggregation is to look at the "loccdc" column and scroll until it is blank. The unique orgIDs will then begin to list in ascending order. Once you see the list go through the highest orgID and start over at the smallest orgID, that is where the unduplicated facility list ends. You will also notice that once the list of orgIDs starts over, the column "locationtype" will begin to have data as well. *Tip: the columns "loccdc" and "locationtype" are **blank** for the rows you want.*
2. Once you identify where the aggregation at orgID starts, click on the first orgID cell and drag until you reach the highest value (before it starts to repeat). Copy the selected cells and paste into a new Excel worksheet or a new sheet within the same worksheet. This is your final list of all unduplicated facilities reporting CLABSI during the timeframe you specified. You will use this list as your facility sampling frame.
3. Generate list of facilities that completed Annual Survey from NHSN:
 - a. On the NHSN landing page, use the Analysis button in the navigation bar and select Reports.
 - b. Use the tree view structure to select Supplemental Reports, Facility-Level Data, and Line Listing – Hospital Survey (2021 and later). If validating a type of facility other than Acute Care Hospitals, select the appropriate corresponding report.
 - c. Select Modify Report and make the following modifications:
 - i. Under the Title/Format tab, select the xls format.
 - ii. Under the Time Period tab, select "completeddate" from the dropdown box, and enter 01/01/2023 for Beginning and 12/31/2023 for Ending. Modify dates as needed for the timeframe being validated.

NOTE: If a facility did not complete the Annual Survey during specified time period, they will not appear in this report.

Modify "Line Listing - Hospital Survey (2021 and later)"

☐ Show descriptive variable names ([Print List](#))

Analysis Data Set: HospSurveyV7

Title/Format

Time Period

Filters

Display Variables

Sort Variables

Display Options

Time Period:

Date Variable

Beginning

Ending

completeddate ▼

01/01/2023

12/31/2023

 Clear Time Period

☐ Enter Date variable/Time period at the time you click the Run button

- iii. Under the Sort Variables tab, double click "surveyYear" in the right-hand box to remove. Find "orgID" in the left-hand box, and double click to move it to the right-hand box.
 - iv. Click the Export button. In the Export Analysis Data Set window, keep the default file type (.csv) and click Export. This will generate a .zip file with a spreadsheet of all facilities that completed the NHSN Annual Survey in the time period designated above.
4. In the facility sampling frame spreadsheet, create a new column for variable "bed size."
 - a. Ensure that facilities are sorted by orgID, in ascending order, in both the facility sampling frame and the Annual Survey line list. Confirm the orgIDs match up before proceeding.
 - b. Copy the numBeds column from the Annual Survey line list and paste into the bed size column in the facility sampling frame spreadsheet. Ensure that the pasted bed size variable is matched to the correct facility.
5. Divide the total facilities in the sampling frame into two strata. Create a new column, "stratum," and assign each facility to either Stratum 1 or Stratum 2:
 - a. Stratum 1: Includes all facilities in the sampling frame that have a bed size of <400.
 - b. Stratum 2: Includes all facilities in the sampling frame that have a bed size of ≥400.
6. Stratum 1:
 - a. If there are 25 or fewer facilities within Stratum 1, select all facilities within Stratum 1 and proceed to Stratum 2.
 - b. If there are more than 25 facilities within Stratum 1, assign a random number to each facility. Sort facilities by random number and select the first 25 facilities.
 - i. Refer to Table 1 below for three methods for random number assignment.
7. Stratum 2:
 - a. If there are 5 or fewer facilities within Stratum 2, select all facilities within Stratum 2 then return to Stratum 1. Select additional facilities from Stratum 1 in descending order, starting with the first facility on the list that was not sampled during step 6, to reach a total of 30 facilities selected.

- b. If there are more than 5 facilities within Stratum 2, assign a random number to each facility. Sort facilities by random number and select the first 5 facilities.
 - i. Refer to Table 1 below for three methods for random number assignment.
 - c. If Stratum 1 has fewer than 25 facilities, return to Stratum 2.
 - 8. Select additional facilities from Stratum 2 in descending order, starting with the first facility on the list not previously sampled, to reach a total of 30 facilities selected.

Table 1. Random number assignment methods	
Option 1: Excel	<ol style="list-style-type: none"> 1. Using the facility list created above, or an HAI line list, insert the command <code>=ROUND(RAND()*1000000,0)</code> into column B and drag to paste this command for each row of the facility list. This will generate a random number for each orgID. 2. Select and copy the values from column B and use the Paste Special (Paste Values) feature to paste the number values into column C. Note: any edit made to the Excel sheet will cause the numbers in column B to recalculate. This is normal and can be ignored if you have an iteration copied. 3. Delete column B so the columns shift left and column C becomes column B. 4. Sort by column B, making sure column A is included in the sort (click on “Expand selection” if a dialog box appears). This is your final list that has been assigned and sorted by a random number.
Option 2: Random Number Generator Website + Excel	<ol style="list-style-type: none"> 1. Identify the total number of facilities from the list created above, or the number of records on HAI line list, 2. Go to https://www.random.org/sequences/ 3. Input 1 as the smallest value, and the total number of facilities/records as the largest value, and click “Get Sequence” 4. Copy the sequence created and paste it into column B of your spreadsheet. 5. Sort by column B, making sure column A is included in the sort (click on “Expand selection” if a dialog box appears). This is your final list that has been assigned and sorted by a random number.
Option 3: SAS Codes	<ol style="list-style-type: none"> 1. Enter the appropriate file path where prompted in the code 2. For medical record random number generation, determine if you need/want the program to create an ‘EoC’ number. If yes, run code as written. If no, delete the lines of code as specified in the program, then run code. 3. The final list, assigned and sorted by a random number, will be exported to the same folder specified in step 1.

Section 2. Download (“freeze”) the facility’s reported data from NHSN

Prior to selecting the medical records sample, use NHSN Analysis Reports and the modifications described below to “freeze” (take a snapshot of) the data and export the facility’s reported CLABSI events. Freeze the data for each facility selected for validation. While in the NHSN application, this would be an opportune time to download each facility’s NHSN Annual Survey, which will be needed during the on-site, or virtual, visit. The Annual Survey will be used to review risk adjustment variables (teaching hospital affiliation, bed count, number of patient days, and number of admissions).

To “freeze” data, select the Analysis tab in the left-hand navigation bar, and then Reports. Select HAI Detailed Reports (Line Lists, Rate Tables, etc.), Device-Associated (DA) Module, CLABSI, and then “Line Listing – All CLABSI Events,” and then click Modify Report.

Suggested Modifications:

- Under the Title/Format tab, select xls as the format. You may also change the title of the report (i.e. <Facility ID> <Freeze Date> NHSN CLABSI Events Line List).
- Under the Time Period tab, go to the Date Variable drop-down and select “eventDateYr.” For both Beginning and Ending, enter 2023, or the year of data to be validated.
- Under the Filters tab, in the row of drop-down boxes, select “orgID,” “equal,” and enter the facility’s orgID number.
 - Optional: Export single report with all facilities, sort by “orgID,” and copy/paste each facility’s data into its own spreadsheet. Save each line list in a secure location.
- Optional: Under the Sort Variables tab, select “eventDate.”
- Click on the “Export” button. Keep the format as-is (.csv) and select the “Export Analysis Data Set using Modifications” radio button. This will generate the line listing in Excel.
- Save the line listing to a secure location. It will be needed again for the medical record selection process in Section 5.

To find a facility’s NHSN Annual Survey, log into NHSN and select Surveys in the Navigation bar, then Find. In the Survey Type drop down menu, select the survey for the type of facility you are validating (for example, FACSRV-PS – Hospital Survey Data, for validation of Acute Care Hospitals). Then, select the appropriate survey year you are validating. Finally, click on the Find button, and a list of facilities and their annual surveys will be generated.

NOTE: Use the **Analysis** button on the Navigation bar and select Reports to export the data. For more information about how to make modifications to these output options, read “How to Modify a Report” found in the Analysis Quick Reference Guide library at: <http://www.cdc.gov/nhsn/PS-Analysis-resources/reference-guides.html>.

Section 3. Notify facilities of the planned validation and request the required laboratory line listings

Suggestions on what information should be included in any outreach to facilities notifying them of their selection can be found in section 2.4 of the 2023 PS EVT. Template letters with suggested format of line listings are located in Appendix 1.

Section 4. Develop the medical record sampling frame for each selected facility

CLABSI in surveillance locations

For CLABSI, the sampling frame is derived from positive laboratory (blood culture) line listings in surveillance locations (SLs). From each selected facility, obtain a complete list of positive blood cultures (PBCs) collected from SLs in 2023 (includes all PBCs taken during SL stay, the day of transfer from the SL, and the day following transfer or discharge) to select the medical record sample before the site visit. NHSN encourages facilities to develop capacity to generate these lists electronically as recurring need for this task is expected, and the creation of manual line listings present an excessive burden.

Note: The term “surveillance locations,” abbreviated SL, is used in the toolkit to indicate that in plan NHSN-reporting locations will be targeted for the validation efforts for CLABSI.

- Validation of CLABSI in NHSN-reporting surveillance locations includes neonatal intensive care unit (NICU) locations

For SL PBCs, identifying organism(s), the MRN, admission date, laboratory specimen number, date of specimen collection (not date of report), resulting first organism genus and species, surveillance location, patient sex, and patient date of birth are required. Additional patient identifiers such as patient name may be helpful. If needed, ask the IP to translate specific patient location information on the laboratory line listings to mapped NHSN-reporting SLs, and ensure that results for all SLs are included. Be sure it is possible to distinguish NICU from adult/pediatric SLs on this line listing to stratify the CLABSI sample. Information about central line use should not be requested; validators will screen for this information while reviewing records. See example of line list template in Appendix 1.2 of the 2023 PS EVT.

Section 5. Medical Record Selection

Use the securely transmitted line listing of PBCs obtained from each selected facility in the following medical record selection process:

1. For each facility, assign a random number to every PBC following steps outlined in Table 1.
2. Sort the list of PBCs by MRN and admission date to generate clusters of PBCs with the same MRN and admission date, also called unique episodes of care (EoC). Create an EoC column where the first EoC = 1. All PBCs in an episode should have the same EoC number.
3. Identify NHSN-reported CLABSI on the facility-provided PBC line listing and assign strata:
 - a. Reference the NHSN CLABSI Events line list created in [Section 2](#) to identify any PBCs reported to NHSN as a CLABSI. Create a new column, “stratum,” and assign any PBCs reported to NHSN as a CLABSI to stratum 1. Additional PBCs in the same EoC as a reported CLABSI should be assigned to stratum 2.
 - b. If any reported CLABSIs, identified from the NHSN CLABSI Events line listing, are missing from the facility-provided PBC line listing, the facility list may be incomplete. Work with the facility to address the missing CLABSIs. Update the facility-provided PBC line list with the missing CLABSI events once the issue has been rectified.
4. Select a simple random sample of 20 reported CLABSIs in NHSN-reporting surveillance locations for review:
 - a. Filter to where stratum = 1.
 - b. Sort by random number.
 - c. Select the first 20 random numbers with a unique EoC number.
 - d. If there are less than 20 reported events, review all stratum 1 and supplement the difference from strata 3 or 4.
5. Identify unreported candidate CLABSI events and stratify by targeted pathogens:
 - a. Filter to where stratum does not equal 1 or 2 and review each PBC to determine if the identified organism is on targeted pathogens list (listed below).
 - b. Sort the unassigned PBCs by identified organism. If more than one organism was identified in an EoC, use the first organism listed.
 - c. If the first listed organism is a targeted pathogen, assign the PBC to stratum 3. If the organism is not a targeted pathogen, assign the PBC to stratum 4.
 - i. Targeted Pathogens**:
 1. *Candida spp.* (yeast)
 2. *Enterococcus spp.*
 3. *Staphylococcus aureus* (includes MRSA, MSSA)
 4. Coagulase-negative *Staphylococcus* (includes most *Staphylococcus spp.* Other than *S. aureus*, MRSA, MSSA)
 5. *Klebsiella spp.*, *E. coli*, or *Pseudomonas spp.* (common gram negatives)
6. Among strata 3 and 4, use location information to identify NICU vs other SL PBCs. Create a new variable, “NICU,” and assign a NICU status (yes/no) to PBC as appropriate. Note: If facility has no NICU, skip to step 8 below and select 5 additional medical records from other SLs for screening sample.
7. Select a targeted NICU screening sample:
 - a. Filter to where NICU = yes and stratum = 3 (targeted pathogens).
 - b. Sort by random number.
 - c. Select the first 5 random numbers with unique EoC numbers.

- d. If 5 PBCs where NICU = yes and stratum = 3 are not available, supplement the difference to reach 5 by using PBCs where NICU = yes and stratum = 4, taking the lowest random number(s) with unique EoCs.
8. Select a targeted non-NICU screening sample:
 - a. Filter to where NICU = no and stratum = 3 (targeted pathogens).
 - b. Sort by random number.
 - c. Select the first 15 random numbers with unique EoC numbers.
 - d. If 15 PBCs where NICU = no and stratum = 3 are not available, supplement to reach 15 by using PBCs NICU = no and stratum = 4, taking the lowest random number(s) with unique EoCs.
9. Review selected PBCs to ensure no EoC are repeated/duplicated. If any EoCs are duplicated, keep the first sampled PBC, and replace the subsequent samples from the pertinent screening sample lists.
10. The final screening sample should contain 20 PBCs with reported CLABSI and 20 candidate PBCs divided among NICU, if available, and non-NICU SLs. If final sample contains less than 40 records, randomly select additional records to reach 40, as possible.
11. If medical records are not well balanced among different targeted pathogens, consider post-selection adjustment to include a variety of these organisms to evaluate a variety of surveillance skills, as noted below.
12. Request the selected medical records in advance of the facility site visit using the template letter found in Appendix 1.3 in the 2023 PS EVT.

*Why Target CLABSI Pathogens?**

The targeted pathogens provide an opportunity to assess a facility's competency in correctly using different components of the NHSN CLABSI definition. For example:

- *Candida* BSI is common in ICU patients receiving parenteral nutrition. Reviewing medical records with *Candida* BSI may provide an opportunity to look for misclassification. *Candida species* and yeast spp. are commonly seen in sputum samples, but infrequently cause true healthcare-associated pneumonia, therefore are considered excluded organisms for PNU criteria, with some exceptions. NHSN cautions against reporting *Candida* pneumonia in immunocompetent patients unless there is evidence of invasive infection on lung biopsy or in pleural fluid under the definitions for PNU. These restrictions are further codified (as prohibitions) under ventilator-associated event (VAE).
- Some facilities that do MRSA active surveillance testing on admission incorrectly assume that MRSA colonization on admission means that a MRSA bloodstream infection would not need to be reviewed for CLABSI.
- Including enteric organisms such as *Enterococcus* and gram-negative rods can demonstrate a facility's ability to distinguish primary bloodstream infection vs. an alternative primary infection like UTI, GIT, or IAB with secondary bloodstream infection. Interested states can also assess use of the mucosal barrier injury reporting definitions, although these are not included in the Toolkit.
- Facilities need to know how to correctly report single and confirmed isolates of common commensal organisms like coagulase-negative *Staphylococcus* and should be able to recognize synonyms (for example *Staphylococcus epidermidis*), used by the microbiology laboratory.

Section 6. Site Visit Activities

6.1 Structured Medical Records Review

Validator blinding and consultation at the facility site-visit

Validator blinding as to HAI status is recommended, when feasible. This can be accomplished by mixing and reviewing the selected medical records before determining which have been reported to NHSN with HAIs.

Medical records should be reviewed in a blinded manner using the 2023 Medical Records Abstraction Tool (MRAT), which can be found at [2023 PSC Data Validation Resources | NHSN | CDC](#), Resources by HAI, CLABSI. This tool includes algorithms and logic designed to establish presence or absence of required criteria for case definitions and provide support to avoid common errors.

For CLABSI validation, when consideration is given to an alternative primary site infection leading to secondary bloodstream infection, use of an appropriate NHSN checklist (available at <https://www.cdc.gov/nhsn/hai-checklists/index.html>) is recommended. These checklists provide a structure to record required elements from the NHSN Patient Safety Component Manual's Chapter 17 criteria. Be sure the selected version is for 2023 definitions.

6.2 Review risk adjustment variables

Have a copy of the facility NHSN Annual Survey available and review surveillance location mapping, location bed size, and teaching hospital status with the IP. A list of CDC locations and descriptions can be found in NHSN Patient Safety Manual Chapter 15.

Review NHSN definitions for teaching hospital types (under Key Terms, Patient Safety Manual Chapter 16), and ensure that facility teaching hospital status is accurate in the NHSN Annual Survey.

6.3 Review denominator collection methods and documentation

Electronically collected CLABSI denominators

If the facility uses electronic denominator data collection, obtain documentation of their denominator validation process and any periodic spot checks. NHSN specifies that electronic denominator counts should fall within 5% of manual counts for three consecutive months before electronic counts can be used. This may be examined pre- or post-visit.

If documentation of electronic denominator validation is not available, the facility should resume manual counting (and ensure staff training) to re-validate electronic counts, and to retain evidence of valid electronic counting (within 5% for 3 months). Facilities should conduct periodic spot checks even after formal validation to prevent lost information due to changing medical records systems or other disruptions.

Manual CLABSI denominator counting methods

Denominator data collection surveys found in sections 6.5 through 6.7 may be administered to the IP contact before or during the site visit. If the facility is manually collecting denominator data, it may be impractical to interview multiple denominator data collectors during the site visit. In this case, collecting contact information during the site visit may be advisable for subsequent administration of surveys by telephone or virtual means. This allows time with the facility to be used efficiently and accommodates interviews with individuals who may work at other times (for example the night shift).

Knowledge of definitions and counting methods is important even in facilities where denominators are reported electronically in order that spot-checks can be conducted periodically. A form for facilities to document required internal validation of electronic denominator counting is provided in Section 6.5.

Facilities may have already administered denominator counting surveys for internal validation purposes. If this is the case, validators may choose to accept their evidence or conduct this survey among a more limited sample of denominator counters.

CLABSI denominator record documentation

While visiting, request original records of denominator data collection paperwork, which can provide insight into the frequency, reliability, and consistency of this task and how omissions are handled. NHSN provides guidance for missing device-associated denominator data that can be found at <https://www.cdc.gov/nhsn/pdfs/gen-support/MissingDenomData-508.pdf>.

Consider whether patient days and central-line days data appear as anticipated when manually counted each day: different ink, different but similar numbers. Determine for what percent of day's data are missing and what was done for reporting on those days. This data is best assessed on site, if possible.

6.4 (Optional) Template for Central Line-Associated Bloodstream Infection (CLABSI) Validation Discrepancies Discussion with Facilities

Please feel free to adapt these templates to meet your jurisdiction's needs to discuss discordant outcomes and request changes

Instructions: For each CLABSI Event with a discordant outcome between facility reporters and validators, record the following (first row: enter facility report; second row: enter validator's recommended changes). Use the Comment area to document reasons for discrepancy, for example: overlooked candidate culture, confusion regarding common commensals, did not meet alternative primary definition, etc. Many states have examined this type of data to identify common errors and direct future education and training. Keep a copy for your records and leave a copy with the facility. F=Facility; V=validator

Pt. ID		Positive blood culture event: first culture date	Select One:			Event date (if LCBI)	If LCBI, MBI* LCBI?	POA, HAI or neither	Central line >2d (y/n)	Location of attribution	CLABSI in Surveillance Locations (y/n)
			Not candidate CLABSI	Alternative primary (specify)	LCBI1, LCBI2, LCBI3*		MBI Yes or MBI No				
	F										
	V										
Comment:											
	F										
	V										
Comment:											
	F										
	V										
Comment:											
	F										
	V										
Comment:											
*LCBI 1, 2, 3 (NHSN): types of laboratory-confirmed bloodstream infection. MBI-LCBI (NHSN) mucosal barrier injury LCBI. See definitions in NHSN Patient Safety Manual Chapter 4.											

6.5 Documentation of Electronic CLABSI Denominator Validation

OrgID/Name of Hospital: _____ Date of Survey: _____

Instructions: NHSN requires that the monthly electronic denominator count falls within a 5% tolerance interval of the monthly manual denominator count for 3 consecutive months before reporting electronic denominator counts for CLABSI. This validation is not conducted during the external survey. The facility is expected to have a copy of this internal validation comparing manual counts to electronic counts available for the validator to review. *If there is no electronic denominator counting at this facility, skip this survey.*
If electronic device denominator counting is used for reporting at this facility, document the NHSN-required validation results below:

Initial electronic denominator validation (when electronic denominator reporting began):

Location name:		Manual count	*Calculated 5% tolerance interval	Electronic count
Month/year:	Patient days			
	Central line days			
Location name:				
Month/year:	Patient days			
	Central line days			
Location name:				
Month/year:	Patient days			
	Central line days			

If available, please document additional information for any more recent electronic denominator validation:

Location name:		Manual count	*Calculated 5% tolerance interval	Electronic count
Month/year	Patient days			
	Central line days			
Location name:				
Month/year	Patient days			
	Central line days			
Location name:				
Month/year:	Patient days			
	Central line days			

*Equation for calculating 5% tolerance interval is: manual count \pm (manual count * 0.05).

Example calculations where manual count = 164 and electronic count = 178:

Eligible 5% tolerance interval = $[164 \pm (164 * 0.05)] = 155.8$ to 172.2

Electronic count 178 falls outside the tolerance interval.

6.6 Contact Information for Manual CLABSI Denominator Validation

Please feel free to adapt this template to meet your jurisdiction's needs

NOTE: If facility ensures annual training updates for denominator counters, and three or more denominator counters show proficiency on the survey, or if facility has already internally surveyed denominator counter proficiency, this can serve as evidence of proficiency.

OrgID / Name of Hospital _____ Date of Survey _____

ID	Name of data collection professional	Surveillance locations covered	Work hours/ Preferred time for telephone survey	Phone number(s)	Supervisor
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					
Add rows as needed					

6.7 CLABSI Denominator Counting Survey (with Key)

<i>Instructions: Administer in person or by telephone, directly to individuals responsible for denominator counting. This form is divided into 2 sections for facilities where these tasks are performed by different persons. The first section contains questions applicable to collecting PATIENT DAYS (questions 1-9). The second section contains questions applicable to collecting CENTRAL LINE DAYS (questions 10-22).</i>				
Facility OrgID:	Name/ID of individual interviewed:	Position: <input type="checkbox"/> IP <input type="checkbox"/> Clerical <input type="checkbox"/> Nursing <input type="checkbox"/> Other (explain)	Interviewer initials:	Date of survey:
CLABSI Denominators		NHSN location(s) covered:		
PATIENT DAYS (for CLABSI denominator counters)			Answer Key:	
1. How are patient days usually collected? (choose one)			<i>If using weekly: Once weekly sampling of denominator data to generate estimated central line days, may be used as an alternative to daily collection in non-oncology ICUs and wards. Sampling may not be used in SCA/ONC locations or NICUs. During the month, the number of patients in the location (patient-days) and the number of patients with at least one central line of any type (central line days) is collected on a designated day each week (for example, every Tuesday), and at the same time each day. The average number of device days per month must be greater than or equal to 75 device days if using weekly denominator collection method. </i>	
	Electronically (document the software system utilized and skip to Q8):			
	Manually (daily/weekly)			
	Some units electronic and some units manual			
	Comment:			
2. Is there a specified time when the denominator count is taken?		<input type="checkbox"/> Yes <input type="checkbox"/> No		The answer should be Yes

3. When is it done?		Counts should be done at a specific time daily, preferably at nearly the same time throughout the facility to avoid errors when patients transfer
4. Describe the method used to count patient days :		From NHSN: Denominator data (patient days and central line days) should be collected at the same time, every day, for each location performing surveillance to ensure that differing collection methods don't inadvertently result in central line days being > patient days.
	Count the number of <u>patients</u> assigned to a unit bed <u>at the same time central line counts are conducted</u>	
	Other (specify):	
5. When reporting monthly patient day total, what is done if there are missing patient day data? (choose one)		NHSN issued specific guidance on imputing values for missing device-associated denominator data https://www.cdc.gov/nhsn/pdfs/gen-support/MissingDenomData-508.pdf
	Report the sum of available daily counts with no adjustment for missing data	
	Estimate or re-create missing data from existing information using our own methods	
	Impute missing values using recent CDC/NHSN guidance	
	Other (specify):	
6. Which best describes your training for denominator (patient days and central line days) counting? (select all that apply)		
	No specific training was provided	Formal training by NHSN or NHSN-trained IP is recommended due to technical aspects of definitions (for example, central line, permanent line, temporary line) and methods (for example, when to count lines, how many to count).
	Peer training (person who previously counted explained their approach to new staff)	
	Formal training by IP	
	Formal training by NHSN (for example, online training)	
	Annual training updates	
	Other (describe):	

7. Which staff member counts patient days and central line days when the “regular” data collector(s) is/are not working?	<input type="checkbox"/> IP <input type="checkbox"/> Another trained counter <input type="checkbox"/> Nobody <input type="checkbox"/> Other (specify)
8. Does your facility have a mechanism in place for quality control of denominator data? <i>(Select one):</i>	
<i>(Electronic data)</i> Yes, data submitted electronically is periodically checked using manual methods	
<i>(Manual data)</i> Yes, manually collected data are periodically counted by more than one staff member	
Yes, other <i>(explain)</i>	
No formal quality control process	
9. Which staff member(s) is/are responsible for entering surveillance locations patient days and central line day data into NHSN?	<input type="checkbox"/> IP <input type="checkbox"/> Counter <input type="checkbox"/> Clerical <input type="checkbox"/> Other (specify)

CENTRAL LINE DAYS <i>(for CLABSI denominator counters only)</i>		Answer Key/Rational
10. How are central line days collected for the unit(s) you oversee? <i>(choose one)</i>		
Electronically (specify <i>software system utilized and skip to Q13</i>):		
Manually (daily/weekly)		
Some units electronic and some units manual		
Comment:		

11. Identify the method used to count central line days : (choose one)		A daily count of <i>the number of patients with a central line in the patient care location during a time period, which is summed for the monthly total</i>
	Count the number of patients with at least one central line at the time surveillance rounds are conducted	
	Count the number of central lines that are in place at the time surveillance rounds are conducted	
	Count the number of central lines that are in use at the time surveillance rounds are conducted	
	Other (specify):	
12. When reporting monthly central line day total, what is done if there are missing central line day data? (choose one)		NHSN issued specific guidance on imputing values for missing device-associated denominator data https://www.cdc.gov/nhsn/pdfs/gen-support/MissingDenomData-508.pdf
	Report the sum of available daily counts with no adjustment for missing data	
	Estimate or re-create missing data using existing information (for example, medical records), then sum	
	Impute missing values using recent CDC/NHSN guidance for missing denominator data	
13. A patient has a radial arterial line and a peripheral IV. How many central line days are counted for this patient on this day?		Zero. The radial arterial line and peripheral IV are not central lines.
14. A patient has a temporary central line and a permanent central line that have both been used during this hospitalization. How many denominator device line days are counted for this patient on this day?		One. Although the patient has two central lines, a device day is defined as the number of patients who have the device, not the number of devices.
15. The patient above with the temporary central line and the permanent central line is on an oncology ward. Should you report one temporary line day, one permanent line day, or both a temporary and a permanent line day?		When a patient in an oncology location has both temporary and permanent lines, the line day is reported as a temporary line day. (This information is detailed in the NHSN PSC Manual, Instructions for Form 57.117I)
16. A patient has a port-a-cath that has not been accessed during this hospital stay, and a peripheral IV that is in use. How many denominator device days are counted for this patient on this day?		One. Beginning in January 2023, central lines that are present on admission should be included in denominator device day counts beginning on the day of admission to an inpatient location. This is regardless of access of the central line. The peripheral IV is not a central line.

17. A port-a-cath was inserted during this admission for planned chemotherapy. It is not in use. How many denominator device days are counted for this patient on this day?	<i>One. If a central line was accessed via placement in an inpatient location during the current admission, it is counted in the denominator device day count each day that it remains in place, whether in use or not.</i>
18. A patient has a central line that was accessed for a blood draw in the ICU yesterday but is not currently in use, and a peripheral IV that is in use. How many central line days are counted for this patient on this day?	<i>One. The central line was accessed in an inpatient location during this stay. All central lines should be included in denominator device day counts once the patient locates to an inpatient location. This is regardless of access of the central line.</i>
19. A patient has a central line that was accessed once for a blood draw in the ED during evaluation leading to admission. The patient is now admitted to an inpatient location, but the line is not currently in use. How many denominator device days are counted for this patient on this day?	<i>One. All central lines should be included in denominator device day counts once the patient locates to an inpatient location. This is regardless of access of the central line.</i>
20. If a central line is removed at 2PM and replaced at 8PM. The central line day count is done at 5PM, should the line be counted?	<div data-bbox="785 675 1129 711"> <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Unknown </div> <div data-bbox="1184 620 1986 740"><i>No. Central line must be in place at time of count.</i></div>

NICU-Specific Central Line Questions <i>(Optional: Check here and skip section if NICU questions do not apply to your job)</i> <input type="checkbox"/>		
21. When reporting central line (CL) days, in neonates, which neonatal weight is used for reporting? <i>(select one)</i>	<input type="checkbox"/> Birth weight <input type="checkbox"/> Current weight	<i>Birth weight</i>
22. Neonates with both a CL and an umbilical catheter (UC) are included in the daily count as: <i>(select one)</i>	<input type="checkbox"/> UC only <input type="checkbox"/> CL only <input type="checkbox"/> 2 separate lines	<i>CL only. No separate reporting of UCs; UCs are considered CLs. Although the patient has two central lines, a device day is defined as the number of patients who have the device, not the number of devices. When reporting the central line, it should be stratified by birth weight.</i>

6.8 (Optional) 2023 CLABSI Validation Summary

*required **conditionally required

Facility Validation Overview	
*Facility ID:	
*Facility Type:	<input type="checkbox"/> Acute care hospital <input type="checkbox"/> Long term acute care hospital (LTAC/LTACH) <input type="checkbox"/> Oncology hospital
*Facility sampling method:	<input type="checkbox"/> CDC Method 1 (Targeted Sampling) <input type="checkbox"/> CDC Method 2 (Cumulative attributable difference) <input type="checkbox"/> CDC Method 3 (Stratified Random Sampling)
Reason Facility was Sampled:	<input type="checkbox"/> All facilities were validated <input type="checkbox"/> Targeted facility (Methods 1 or 2) <input type="checkbox"/> Randomly selected

Numerator Validation			
*Sampling information for numerator validation at this facility:			
Event	Sampling Frame Elements	Sampling Frame (# episodes/procedures eligible for review during timeframe)	Total # events facility reported to NHSN for timeframe (before validation)
CLABSI (including NICU)	Medical records with PBC(s)	_____	_____
*CLABSI in surveillance locations (including NICU) Validation Results:			
Event Determination	Validation: Yes - CLABSI	Validation: No – Not CLABSI	
Facility: Yes - Date-matched CLABSI reported	a. _____	b. _____	
Facility: No - Date-matched CLABSI NOT reported	c. _____	d. _____	

Denominator Validation: Central Line and Patient days for CLABSI	
**Which method was used by this facility for CLABSI in surveillance locations denominator (patient days and central line days) counting for this timeframe?	<input type="checkbox"/> Manual counting: <input type="checkbox"/> Daily <input type="checkbox"/> Weekly sampling++ <input type="checkbox"/> Electronic counting <input type="checkbox"/> Both manual and electronic counting
++ Only ICU and ward location types with an average of 75 or more central line-days per month are eligible to use this method.	

**Has this facility completed an internal validation of CLABSI in surveillance locations denominator data for this year?

- ☐ Yes
☐ No

**If yes, provide the following information for all locations and months validated:

Location of validation	Month of validation	Validation method (enter A, B, or C)	Count 1	Count 2

Notes: If Method A is chosen, Count 1 should be "Usual Count" and Count 2 should be "Expert (Referent) Count."
 If Method B is chosen, Count 1 should be "Usual Count" and Count 2 should be "Patient days (Referent) Count."
 If Method C is chosen, Count 1 should be "Manual Count" and Count 2 should be "Electronic Count."

Validation of manual denominator data counting requires either:

- Method A – Concurrent dual counting (with more experienced counter as reference) for \geq three months OR
- Method B – Concurrent patient days data (ADT-Admission/Discharge/Transfer or other reference) and manual counting for \geq three consecutive months

Validation of electronic denominator data counting requires:

- Method C – Concurrent manual denominator counting (reference) vs. electronic data for \geq three months

NHSN Inpatient Location Validation: MAPPING

**Do any inpatient locations require mapping or re-mapping within NHSN?

- ☐ Yes
☐ No

**If yes, indicate which locations need to be mapped/re-mapped and recommendations:

Location	Current CDC location code designation	Current bed count	Recommended CDC location code designation	Recommended bed count

Add rows as needed.

**How does this facility obtain inpatient admissions data?	<input type="checkbox"/> Electronic from billing <input type="checkbox"/> Electronic from vendor system <input type="checkbox"/> Electronic from ADT <input type="checkbox"/> Other (specify): _____
**How does this facility obtain inpatient patient days data?	<input type="checkbox"/> Electronic from billing <input type="checkbox"/> Electronic from vendor system <input type="checkbox"/> Electronic from ADT <input type="checkbox"/> Other (specify): _____

Risk Adjustment Variable Validation	
**CLABSI Surveillance location (SL) mapping (including NICUs)	
Number of SLs correctly mapped in NHSN (including NICUs):	
Number of SLs incorrectly mapped (including NICUs):	
Number of SLs (including NICUs) omitted from mapping:	
Number of SLs mapping errors:	
**Teaching hospital affiliation (CLABSI in surveillance locations)	
Facility teaching hospital affiliation reported on 2023 NHSN annual facility survey:	<input type="checkbox"/> Non-teaching <input type="checkbox"/> Major <input type="checkbox"/> Graduate <input type="checkbox"/> Undergraduate <input type="checkbox"/> N/A (LTACH)
Is facility teaching hospital affiliation correct?	<input type="checkbox"/> Yes <input type="checkbox"/> No

Comments